

Markscheme

May 2019

Physics

Higher level

Paper 2

15 pages

No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without written permission from the IB.

Additionally, the license tied with this product prohibits commercial use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, is not permitted and is subject to the IB's prior written consent via a license. More information on how to request a license can be obtained from <http://www.ibo.org/contact-the-ib/media-inquiries/for-publishers/guidance-for-third-party-publishers-and-providers/how-to-apply-for-a-license>.

Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite de l'IB.

De plus, la licence associée à ce produit interdit toute utilisation commerciale de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, n'est pas autorisée et est soumise au consentement écrit préalable de l'IB par l'intermédiaire d'une licence. Pour plus d'informations sur la procédure à suivre pour demander une licence, rendez-vous à l'adresse <http://www.ibo.org/fr/contact-the-ib/media-inquiries/for-publishers/guidance-for-third-party-publishers-and-providers/how-to-apply-for-a-license>.

No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin que medie la autorización escrita del IB.

Además, la licencia vinculada a este producto prohíbe el uso con fines comerciales de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales— no está permitido y estará sujeto al otorgamiento previo de una licencia escrita por parte del IB. En este enlace encontrará más información sobre cómo solicitar una licencia: <http://www.ibo.org/es/contact-the-ib/media-inquiries/for-publishers/guidance-for-third-party-publishers-and-providers/how-to-apply-for-a-license>.

Subject Details: Physics HL Paper 2 Markscheme**Mark Allocation**

Candidates are required to answer ALL questions. Maximum total = [90 marks].

1. Each row in the “Question” column relates to the smallest subpart of the question.
2. The maximum mark for each question subpart is indicated in the “Total” column.
3. Each marking point in the “Answers” column is shown by means of a tick (✓) at the end of the marking point.
4. A question subpart may have more marking points than the total allows. This will be indicated by “max” written after the mark in the “Total” column. The related rubric, if necessary, will be outlined in the “Notes” column.
5. An alternative wording is indicated in the “Answers” column by a slash (/). Either wording can be accepted.
6. An alternative answer is indicated in the “Answers” column by “**OR**” between the alternatives. Either answer can be accepted.
7. Words in angled brackets « » in the “Answers” column are not necessary to gain the mark.
8. Words that are underlined are essential for the mark.
9. The order of marking points does not have to be as in the “Answers” column, unless stated otherwise in the “Notes” column.
10. If the candidate’s answer has the same “meaning” or can be clearly interpreted as being of equivalent significance, detail and validity as that in the “Answers” column then award the mark. Where this point is considered to be particularly relevant in a question it is emphasized by **OWTTE** (or words to that effect) in the “Notes” column.
11. Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
12. Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then **follow through** marks should be awarded. When marking, indicate this by adding **ECF** (error carried forward) on the script. “Allow ECF” will be displayed in the “Notes” column.
13. Do **not** penalize candidates for errors in units or significant figures, **unless** it is specifically referred to in the “Notes” column.
14. Allow reasonable substitutions where in common usage, eg ° for rad.

| Question | | | Answers | Notes | Total |
|----------|---|-----|--|--|-------|
| 1. | a | i | time taken $\frac{2.0 \times 10^4}{7}$ «= 2860 s» = 2900«s» ✓ | <i>Must see at least two s.f.</i> | 1 |
| 1. | a | ii | use of $E = qV$ OR energy = $4.3 \times 10^4 \times 16$ «= 6.88×10^5 J» ✓ power = 241 «W» ✓ | <i>Accept 229 W – 241 W depending on the exact value of t used from ai.</i> <i>Must see at least three s.f.</i> | 2 |
| 1. | a | iii | use of power = force x speed OR force x distance = power x time ✓ 34 «N» ✓ | <i>Accept 34 N – 36 N.</i> | 2 |
| 1. | b | i | $66 \text{ g} \sin(3^\circ) = 34$ «N» ✓ | | 1 |
| 1. | b | ii | total force $34 + 34 = 68$ «N» ✓ 3.5 «ms ⁻¹ » ✓ | <i>Look for ECF from aiii and bi.</i> <i>Accept 3.4 – 3.5 «ms⁻¹».</i> <i>Award [0] for solutions involving use of KE.</i> | 2 |

(continued...)

(Question 1 continued)

| Question | | | Answers | Notes | Total |
|----------|---|----|--|---------------------|-------|
| 1. | c | | «maximum» distance will decrease OWTTE ✓ because opposing/resistive force has increased OR because more energy is transferred to GPE OR because velocity has decreased OR increased mass means more work required «to move up the hill» ✓ | | 2 |
| 1. | d | | 4 V dropped across battery OR $R_{\text{circuit}} = 1.85 \Omega$ ✓ so internal resistance $= \frac{4.0}{6.5} = 0.62 \text{ «}\Omega\text{»}$ ✓ | | 2 |
| 1. | e | i | $\frac{16}{5} = 3.2 \text{ «V»}$ ✓ | | 1 |
| 1. | e | ii | ALTERNATIVE 1: $2.5r = 0.62$ ✓ $r = 0.25 \text{ «}\Omega\text{»}$ ✓ ALTERNATIVE 2: $\frac{0.62}{5} = 0.124 \text{ «}\Omega\text{»}$ ✓ $r = 2(0.124) = 0.248 \text{ «}\Omega\text{»}$ ✓ | Allow ECF from (d). | 2 |

| Question | | | Answers | Notes | Total |
|----------|---|-----|--|--|-------|
| 2. | a | i | read off between 17 and 19 «deg» ✓ correct use of $d = \frac{\lambda}{\sin\theta} = 7.8 \times 10^{-15}$ «m» ✓ so radius = $\frac{7.8}{2}$ «fm» = 3.9 «fm» ✓ | Award ecf for wrong angle in MP1. Answer for MP3 must show at least 2 sf. | 3 |
| 2. | a | ii | $R_{\text{Th}} = R_{\text{Si}} \left(\frac{A_{\text{Th}}}{A_{\text{Si}}}\right)^{\frac{1}{3}}$ or substitution ✓ 7.4 «fm» ✓ | | 2 |
| 2. | a | iii | electron wavelength shorter than alpha particles (thus increased resolution) OR electron is not subject to strong nuclear force ✓ | | 1 |
| 2. | a | iv | nuclear forces act ✓ nuclear recoil occurs ✓ significant penetration into nucleus / probing internal structure of individual nucleons ✓ incident particles are relativistic ✓ | | 2 max |

(continued...)

(Question 2 continued)

| Question | | | Answers | Notes | Total |
|----------|---|-----|--|-------|-------|
| 2. | b | i | ${}_{15}^{30}\text{P} \rightarrow ({}_{14}^{30}\text{Si}) \checkmark$ $+ {}_{+1}^0\text{e} + \nu_e \checkmark$ | | 2 |
| 2. | b | ii | <p>correct change of either u to d \checkmark W^+ shown \checkmark correct arrow directions for positron and electron neutrino \checkmark</p> | | 3 |
| 2. | b | iii | quarks cannot be directly observed as free particles/must remain bound to other quarks/quarks cannot be isolated \checkmark because energy given to nucleon creates other particles rather than freeing quarks/ OWTTE \checkmark | | 2 |


(continued...)

(Question 2 continued)

| Question | | Answers | Notes | Total |
|----------|---|---|--|-------|
| 2. | c | models need testing/new information may change models/new technology may bring new information/Models can be revised/ OWTTE ✓ | | 1 |
| 3. | a | two waves superpose/mention of superposition/mention of «constructive» interference ✓ they arrive in phase/there is a path length difference of an integer number of wavelengths ✓ | | 2 |
| 3. | b | path difference = 0.062 «m» ✓ so wavelength = 0.031 «m» ✓ frequency = 9.7×10^9 «Hz» ✓ | Award [2 max] for 4.8×10^9 Hz. | 3 |
| 3. | c | intensity is modulated by a single slit diffraction envelope OR intensity varies with distance OR points are different distances from the slits ✓ | | 1 |

(continued...)

(Question 3 continued)

| Question | | Answers | Notes | Total |
|----------|---|---|--|-------|
| 3. | d |  <p>cos² variation shown ✓ with zero at 90° (by eye) ✓</p> | <p>Award [1 max] for an inverted curve with maximum at 90°.</p> | 2 |

| Question | | | Answers | Notes | Total |
|----------|---|-----|---|--|-------|
| 4. | a | | weight of cylinder = $Ahg\rho$ ✓ pressure = $\frac{F}{A} = \frac{Ahg\rho}{A}$ ✓ | | 2 |
| 4. | b | i | use of $PV = nRT$ and $V = \text{Area} \times (0.190)$ seen ✓ substitution of $P = p_0 + p_m$ «re-arrangement to give answer» ✓ | | 2 |
| 4. | b | ii | recognition that $\frac{nRT}{A}$ is constant OR $190p_0 + 190p_m = 208p_0 - 208p_m$ OR $p_0 = \frac{398}{18} p_m$ ✓ pressure due to mercury $p_m = 0.035 \times 1.36 \times 10^4 \times 9.81 (= 4.67 \times 10^3 \text{ Pa})$ ✓ 1.03×10^5 ✓ Pa OR Nm^{-2} OR $\text{kgm}^{-1}\text{s}^{-2}$ ✓ | <i>Award MP4 for any correct unit of pressure (eg "mm of mercury / Hg").</i> | 4 |
| 4. | b | iii | same number of particles to collide with a larger surface area OR greater volume with constant rms speed decreases collision frequency ✓ | | 1 |

| Question | | | Answers | Notes | Total |
|----------|---|----|---|---------------------------------|-------|
| 5. | a | i | gravitational attraction/force/field «of the planet/Mars» ✓ | <i>Do not accept "gravity".</i> | 1 |
| 5. | a | ii | the force/field and the velocity/displacement are at 90° to each other OR there is no change in GPE of the moon/Phobos ✓ | | 1 |
| 5. | b | i | <p>ALTERNATE 1 «using fundamental equations» use of Universal gravitational force/acceleration/orbital velocity equations ✓ equating to centripetal force or acceleration. ✓ rearranges to get $k = \frac{G}{4\pi^2}$ ✓</p> <p>ALTERNATE 2 «starting with $\frac{R^3}{T^2} = kM$» substitution of proper equation for T from orbital motion equations ✓ substitution of proper equation for M OR R from orbital motion equations ✓ rearranges to get $k = \frac{G}{4\pi^2}$ ✓</p> | | 3 |

(continued...)

(Question 5 continued)

| Question | | | Answers | Notes | Total |
|----------|---|----|--|--------------------|-------|
| 5. | b | ii | $m_{\text{Mars}} = \left(\frac{R_{\text{Mars}}}{R_{\text{Earth}}} \right)^3 \left(\frac{T_{\text{Earth}}}{T_{\text{Mars}}} \right)^2 m_{\text{Earth}}$ or other consistent re-arrangement ✓ 6.4×10^{23} «kg» ✓ | | 2 |
| 5. | c | | read off separation at maximum potential 0.9 ✓ equating of gravitational field strength of earth and moon at that location OR $\frac{M_E}{0.9^2} = \frac{M_M}{0.1^2}$ ✓ 7.4×10^{22} «kg» ✓ | Allow ECF from MP1 | 3 |

| Question | | | Answers | Notes | Total |
|----------|---|-----|--|--|-------|
| 6. | a | i | identifies T as 2.25 s ✓ L = 1.26 m ✓ 1.3 / 1.26 «m» ✓ | Accept <u>any</u> number of s.f. for MP2. Accept <u>any</u> answer with 2 <u>or</u> 3 s.f. for MP3. | 3 |
| 6. | a | ii | X labels any point <u>on the curve</u> where $E_k \frac{1}{4}$ of maximum/5 mJ ✓ | | 1 |
| 6. | a | iii | $\frac{1}{2}mv^2 = 20 \times 10^{-3}$ seen OR $\frac{1}{2} \times 7.5 \times 10^{-2} \times v^2 = 20 \times 10^{-3}$ ✓ 0.73 «m s ⁻¹ » ✓ | Must see at least 2 s.f. for MP2. | 2 |
| 6. | b | i | 0.40 «m s ⁻¹ » ✓ | | 1 |
| 6. | b | ii | initial energy 24 mJ and final energy 12 mJ ✓ energy is lost/unequal /change in energy is 12 mJ ✓ inelastic collisions occur when energy is lost ✓ | | 3 |
| 6. | b | iii | graph with same period but inverted ✓ amplitude one half of the original/two boxes throughout (by eye) ✓ | | 2 |
| 6. | b | iv | mention of Doppler effect ✓ there is a change in the wavelength of the reflected wave ✓ because the wave speed is constant, there is a change in frequency ✓ | | 3 |

| Question | | | Answers | Notes | Total |
|----------|---|----|--|--|-------|
| 7. | a | i | $T = \left(\frac{1360}{\sigma} \right)^{0.25} \checkmark$ <p>390 «K» ✓</p> | <p><i>Must see 1360 (from data booklet) used for MP1.</i></p> <p><i>Must see at least 2 s.f.</i></p> | 2 |
| 7. | a | ii | <p>energy/Power/Intensity lower at B ✓</p> <p>connection made between energy/power/intensity and temperature of blackbody ✓</p> | | 2 |
| 7. | b | | <p>(28%) of sun's energy is scattered/reflected by earth's atmosphere OR only 72% of incident energy gets absorbed by blackbody ✓</p> | <p><i>Must be clear that the energy is being scattered by the atmosphere.</i></p> | 1 |

| Question | | | Answers | Notes | Total |
|----------|---|-----|---|-------|-------|
| 8. | a | i | $\text{length} = \frac{d \times C}{\text{width} \times \epsilon} \checkmark$ $= 0.33 \text{ «m»} \checkmark$ so 0.66/0.67 «m» «as two lengths required» \checkmark | | 3 |
| 8. | a | ii | $1.5 \times 10^6 \times 55 \times 10^{-6} = 83 \text{ «V»} \checkmark$ $q \text{ «= CV»} = 5.6 \times 10^{-6} \text{ «C»} \checkmark$ | | 2 |
| 8. | b | i | $0.5 = e^{-\frac{t}{RC}} = e^{-\frac{t}{1200 \times 6.8 \times 10^{-8}}} \checkmark$ $t = \text{«-»} 1200 \times 6.8 \times 10^{-8} \times \ln 0.5 \checkmark$ $5.7 \times 10^{-5} \text{ «s»} \checkmark$ <p>OR</p> use of $t_{\frac{1}{2}} = RC \times \ln 2 \checkmark$ $1200 \times 6.8 \times 10^{-8} \times 0.693 \checkmark$ $5.7 \times 10^{-5} \text{ «s»} \checkmark$ | | 3 |
| 8. | b | ii | mention of Faraday's law \checkmark indicating that changing current in discharge circuit leads to change in flux in coil/change in magnetic field «and induced emf» \checkmark | | 2 |
| 8. | b | iii | decrease/reduce \checkmark resistance (R) OR capacitance (C) \checkmark | | 2 |